For Research Use Only

Phospho-S6 Ribosomal protein (Ser236) Recombinant antibody



Catalog Number:80206-1-RR

Basic Information

Catalog Number: 80206-1-RR

Size: 250 μg/ml Source: Rabbit

GeneID (NCBI): 6194 **UNIPROT ID:** P62753

Full Name: Isotype: ribosomal protein S6

> Calculated MW: 29 kDa Observed MW: 32 kDa

GenBank Accession Number:

BC000524

Purification Method:

Protein A purification

CloneNo.: 7K17

Recommended Dilutions: WB 1:2000-1:10000

Applications

Tested Applications: FC, WB, ELISA Species Specificity: Human, mouse

Positive Controls:

WB: HEK-293 cells, MCF-7 cells, Calyculin A treated HEK-293 cells, IGF-1 treated MCF-7 cells

Background Information

Ribosomal protein S6 (RPS6) is one of the components of the 40S ribosomal subunit. RPS6 has been functionally regarded as the stimulator and/or inhibitor of certain types of mRNA translation, as well as the regulator of cellular metabolisms, cells size, survival and proliferation. RPS6 is phosphorylated at multiple sites, comprised between Ser235 and Ser247, by the p70 rpS6 kinase (S6K) 1, which is a major downstream effector of the mammalian target of rapamycin complex 1 (mTORC1). Phosphorylation of RPS6 at the dual site Ser235/236 occurs also independently of mTORC1, via the p90 ribosomal S6 kinases (RSK), which are activated by the extracellular signal-regulated kinases (ERK). Recent studies performed in pancreatic β-cells identified PKA as an additional RPS6 kinase, specifically involved in the phosphorylation of Ser235/236. (PMID: 26490682, PMID: 21814187, PMID: 31112404). 80206-1-RR specifically recognizes the phosphorylation site of Ser236 or dual site Ser235/236.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Non-treated and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 80206-1-RR (Phospho-56 Ribosomal protein (Ser236) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



1X10^6 HEK-293 cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug Anti-Human Phospho-56 Ribosomal protein (Ser236) (80206-1-RR, Clone:7K17) and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:1000, or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.