For Research Use Only

Vimentin Recombinant antibody

Catalog Number:80232-1-RR

Featured Product



Basic Information

Catalog Number: 80232-1-RR

Size: GeneID (NCBI): 500 μ g/ml 7431

Source: ENSEMBL Gene ID:
Rabbit ENSG0000026025
Isotype: UNIPROT ID:
IgG P08670
Immunogen Catalog Number: Full Name:
AG0489 vimentin

Calculated MW: 466 aa, 54 kDa Observed MW:

Observed MV 50-55 kDa

GenBank Accession Number:

BC000163

Purification Method:

Protein A purification

CloneNo.: 6K21

Recommended Dilutions: WB 1:20000-1:100000 IHC 1:1250-1:5000 IF 1:200-1:800

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity: Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: MCF-7 cells, U-251 cells, A549 cells, HeLa cells, U2OS cells, NIH/3T3 cells, 4T1 cells, ROS1728 cells, HSC-T6 cells

IHC : human breast cancer tissue, human liver cancer tissue

IF: HepG2 cells,

Background Information

Vimentin, also named as VIM, belongs to the intermediate filament family. Vimentin is class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is important for stabilizing the architecture of the cytoplasm. Monocyte-derived macrophages secrete vimentin into the extracellular space in vitro. Secretion of vimentin was enhanced by the proinflammatory cytokine tumor necrosis factor-alpha (TNFA; 191160) and inhibited by the antiinflammatory cytokine IL10 (124092), suggesting that vimentin is involved in the immune response. Vimentin has specialized functions that contribute to specific dynamic cellular processes. As a phosphoprotein, 55-60 kDa of vimentin proteins can be observed due to the different phosphorylation level. Isoforms of vimentin (49 kDa and 60 kDa) had also been reported. (PMID: 8640945, 22728585).


Storage

Storage: Store at -20°C. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 80232-1-RR (Vimentin antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



WB result of Vimentin antibody (80232-1-RR; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Vimentin transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 80232-1-RR (Vimentin antibody) at dilution of 1:2500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using Vimentin antibody (80232-1-RR, Clone: 6K21) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).