For Research Use Only

TNF-alpha Recombinant antibody

Catalog Number:80258-6-RR



Basic Information

Catalog Number: GenBank Accession Number:

 400 ug/ml
 21926

 Source:
 UNIPROT ID:

 Rabbit
 P06804

 Isotype:
 Full Name:

IgG tumor necrosis factor

Observed MW: 25-26 kDa Purification Method:

240702B7

Protein A purfication CloneNo.:

Recommended Dilutions: WB 1:2000-1:10000 IF/ICC 1:125-1:500

Applications

Tested Applications: WB, IF/ICC, ELISA Species Specificity:

mouse

Positive Controls:

WB: PMA, LPS and Brefeldin A treated RAW 264.7 cells, IF/ICC: LPS and Brefeldin A treated RAW 264.7 cells,

Background Information

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. Mouse and human TNF-alpha share 79% amino acid sequence identity. Unlike human TNF-alpha, the mouse form is glycosylated. In mouse deficiency of this gene is associated with defects in response to bacterial infection, with defects in forming organized follicular dendritic cell networks and germinal centers, and with a lack of primary B cell follicles.

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



PMA, LPS and Brefeldin A treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 80258-6-RR (Tnfa antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed LPS and Brefeldin A treated RAW 264.7 cells using Tnfa antibody (80258-6-RR, Clone: 240702B7) at dilution of 1:250 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L) (5A00013-2).



Biolayer interferometry (BLL) kinetic assays of 80258-6-RR against Mouse Tnfa were performed. The affinity constant is below 1 pM.