For Research Use Only

ADRP Recombinant antibody

Catalog Number:80362-2-RR




Basic Information	Catalog Number: 80362-2-RR	GenBank Accession Number: BC005127	Purification Method: Protein A purification		
	Size:	GenelD (NCBI):	CloneNo.:		
	1000 µg/ml	123	230274F4		
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG7539	UNIPROT ID: Q99541 Full Name: adipose differentiation-related protein	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500 IF 1:200-1:800		
				ed 11 1.200 1.000	
				Calculated MW:	
			48 kDa		
		Observed MW: 48 kDa			
		Applications	Tested Applications: WB, IF/ICC, IHC, ELISA		ve Controls:
				WB : T	WB : THP-1 cells, HepG2 cells, K-562 cells
Species Specificity: Human. mouse	IHC : r		nouse liver tissue,		
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			eic acid treated HeLa cells,		
Background Information	ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT family which is responsible for the transportation of lipids and the formation of lipid droplets. ADRP is localized on the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.				
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.			


For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data

temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 80362-2-RR (ADRP antibody) at dilution of 1:5000 incubated at room




Immunofluorescent analysis of (4% PFA) fixed oleic acid treated HeLa cells using ADRP antibody (80362-2-RR, Clone: 230274F4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).

Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 80362-2-RR (ADRP antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffor (pH 0) EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 80362-2-RR (ADRP antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH o) EDTÁ buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 80362-2-RR (ADRP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 80362-2-RR against Human ADRP were performed. The affinity constant is below 1 pM.