## For Research Use Only

## CD31 Recombinant antibody, PBS Only



**Purification Method:** 

CloneNo.:

4N13

Protein A purification

Catalog Number: 80530-1-PBS

**Basic Information** 

Catalog Number:

80530-1-PBS

Size: GeneID (NCBI): 1 mg/ml 5175

Source: ENSEMBL Gene ID:
Rabbit ENSG0000261371
Isotype: UNIPROT ID:

IgG P16284
Immunogen Catalog Number: Full Name:

AG1787 platelet/endothelial cell adhesion

Calculated MW: 83 kDa Observed MW:

GenBank Accession Number:

BC022512

130 kDa

molecule

**Applications** 

Tested Applications: WB,Indirect ELISA,IF Species Specificity: Human, Mouse


## **Background Information**

Platelet endothelial cell adhesion molecule-1 (PECAM-1, CD31) is a member of the immunoglobulin gene superfamily of cell adhesion molecules. CD31 is a transmembrane glycoprotein that is highly expressed on the surface of the endothelium, making up a large portion of its intracellular junctions. PECAM-1 is also present on the surface of hematopoietic cells and immune cells including platelets, monocytes, neutrophils, natural killer cells, megakaryocytes and some types of T-cell (PMID: 9011572). As well as its role in cell-cell adhesion, PECAM-1 functions as a signaling receptor, and is involved in important physiological events such as nitric oxide production, regulation of T-cell immunity and tolerance, leukocyte transendothelial migration and inflammation and angiogenesis (PMID: 21183735; 20978210; 17872453; 20634489).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


## **Selected Validation Data**



Untreated and PNGase F-treated lysates of human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 80530-1-RR (CD31 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80530-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 80530-1-RR (CD31 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80530-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed bEnd.3 cells using CD31 antibody (80530-1-RR, Clone: 4N13) at dilution of 1:500 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Alpha Tubulin antibody (66031-1-Ig, Clone: 1E4C11, red). This data was developed using the same antibody clone with 80530-1-PBS in a different storage buffer formulation.