

For Research Use Only

p38 MAPK Recombinant antibody, PBS Only



Catalog Number:80821-3-PBS

Basic Information

Catalog Number: 80821-3-PBS	GenBank Accession Number: BC031574	Purification Method: Protein A purification
Size: 1 mg/ml	GeneID (NCBI): 1432	CloneNo.: 230263H7
Source: Rabbit	UNIPROT ID: Q16539	
Isotype: IgG	Full Name: mitogen-activated protein kinase 14	
Immunogen Catalog Number: AG5115	Calculated MW: 360 aa, 41 kDa	
	Observed MW: 38-42 kDa	

Applications

Tested Applications:
WB, Indirect ELISA, FC

Species Specificity:
Human, mouse

Background Information

MAPK14 (mitogen-activated protein kinase 14) is also named as SAPIK2A, p38MAPK, CSBP1, RK, p38, EXIP, Mxi2, CSBP2, PRKM14, PRKM15, CSPB1, p38ALPHA and belongs to the MAP kinase subfamily. MAPK14-signaling is a central pathway for the integration of instructive signals in dendritic cells for T(H)17 differentiation and inflammation (PMID:22231518). It plays an important role in the regulation of hematopoietic stem cell self-renewal in vitro and inhibition of MAPK14 activation with a small molecule inhibitor may represent a novel approach to promote ex vivo expansion of hematopoietic stem cell (PMID:21198398). This protein has 4 isoforms produced by alternative splicing.

Storage

Storage:
Store at -80°C.

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 80821-3-RR (p38 MAPK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80821-3-PBS in a different storage buffer formulation.



1x10⁶ HeLa cells were intracellularly stained with 0.25 ug Anti-Human p38 MAPK (80821-3-RR, Clone:230263H7) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set. This data was developed using the same antibody clone with 80821-3-PBS in a different storage buffer formulation.



1x10⁶ Jurkat cells were intracellularly stained with 0.25 ug Anti-Human p38 MAPK (80821-3-RR, Clone:230263H7) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set. This data was developed using the same antibody clone with 80821-3-PBS in a different storage buffer formulation.