

For Research Use Only

Phospho-TBC1D1 (Ser660) Recombinant antibody, PBS Only



Catalog Number: 81363-2-PBS

Basic Information

Catalog Number:

81363-2-PBS

Size:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC050321

GeneID (NCBI):

23216

UNIPROT ID:

Q86T10

Full Name:

TBC1 (tre-2/USP6, BUB2, cdc16)
domain family, member 1

Calculated MW:

133 kDa

Observed MW:

160 kDa

Purification Method:

Protein A purification

CloneNo.:

7N19

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

Human, mouse

Background Information

TBC1D1 belongs to the family of RabGAPs and its physiological roles in glucose homeostasis and its implication in human diseases have already been characterized in skeletal muscle and adipocytes in a large number of independent studies. TBC1D1 including twophospho-tyrosine-binding (PTB) domains, a calmodulin-binding domain (CBD) and the functional GAP domain responsible for catalyzing GTP hydrolysis. TBC1D1 is phosphorylated by AMPK at Ser660 and Thr237, and by AKT at Thr596. The RabGAPs TBC1D1 is important regulators of glucose homeostasis in skeletal muscle and adipose tissue. (PMID: 31627187, PMID: 18477703)

Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 81363-2-RR (Phospho-TBC1D1 (Ser660) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin antibody as loading control. This data was developed using the same antibody clone with 81363-2-PBS in a different storage buffer formulation.



Non-treated and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 81363-2-RR (Phospho-TBC1D1 (Ser660) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin antibody as loading control. This data was developed using the same antibody clone with 81363-2-PBS in a different storage buffer formulation.