

For Research Use Only

SCD Recombinant antibody

Catalog Number: 81468-2-RR



Basic Information

Catalog Number:

81468-2-RR

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG29156

GenBank Accession Number:

BC005807

GeneID (NCBI):

6319

UNIPROT ID:

O00767

Full Name:

stearoyl-CoA desaturase (delta-9-desaturase)

Calculated MW:

355 aa, 41 kDa

Observed MW:

28-42 kDa

Purification Method:

Protein A purification

CloneNo.:

241487B5

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

human

Positive Controls:

WB: A431 cells, HepG2 cells, A375 cells, A549 cells, HEK-293T cells

Background Information

SCD1 (stearoyl-CoA desaturase) is a microsomal fatty acid monodesaturase, which catalyses the committed step in the biosynthesis of mono-unsaturated fatty acids from saturated fatty acids (PMID:10946019). SCD1 and SCD2 are the main isoforms expressed in mouse liver and brain respectively (PMID:15907797). The formation of homodimers and oligomers is an intrinsic property of SCD proteins. SCD1 is a multi-pass membrane protein and detected double bands of 37-42 kDa. The degradation product of 28 kDa may be caused by a major cleavage site at the C-terminus (PMID:15610069, PMID: 9843580).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81468-2-RR (SCD antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1×10^6 HepG2 cells were intracellularly stained with 0.2 μ g SCD Recombinant Antibody (81468-2-RR, Clone:241487B5) and Multi-rAb Coralite[®] Plus 647-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR005)(red), or 0.2 μ g Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1×10^6 MCF-7 cells were intracellularly stained with 0.2 μ g SCD Recombinant Antibody (81468-2-RR, Clone:241487B5) and Multi-rAb Coralite[®] Plus 647-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR005)(red), or 0.2 μ g Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).