

For Research Use Only

# Phospho-CHEK2 (Thr68) Recombinant antibody, PBS Only



Catalog Number: 81740-1-PBS

## Basic Information

<b>Catalog Number:</b> 81740-1-PBS	<b>GenBank Accession Number:</b> BC004207	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 1 mg/ml	<b>GeneID (NCBI):</b> 11200	<b>CloneNo.:</b> 1L2
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> O96017	
<b>Isotype:</b> IgG	<b>Full Name:</b> CHK2 checkpoint homolog (S. pombe)	
	<b>Calculated MW:</b> 61 kDa	
	<b>Observed MW:</b> 65 kDa	

## Applications

**Tested Applications:**  
WB, Indirect ELISA

**Species Specificity:**  
Human

## Background Information

Serine/threonine-protein kinase Chk2 (CHEK2) is a serine/threonine kinase which is activated upon DNA damage and is implicated in pathways that govern DNA repair, cell cycle arrest or apoptosis in response to the initial damage. ATM phosphorylates CHEK2 on T68. Phosphorylation on T68 and subsequent full activation of CHEK2 was shown to require priming phosphorylation on adjacent residues by Polo-like kinase 3 (PLK3) and the dual specificity tyrosine and serine/threonine kinase TTK/hMPS1. Additionally TTK appears to phosphorylate T68. Phosphorylation of T68 promotes the binding of the N-terminal SQ/TQ-rich cluster of one CHEK2 molecule with the FHA domain of another CHEK2 molecule. (PMID: 28553140, PMID: 18004398, PMID: 33322746)

## Storage

**Storage:**  
Store at -80°C.

**Storage Buffer:**  
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Non-treated PC-3 and MMS treated PC-3 cells were subjected to SDS PAGE followed by western blot with 81740-1-RR (Phospho-CHEK2 (Thr68) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin antibody as loading control. This data was developed using the same antibody clone with 81740-1-PBS in a different storage buffer formulation.



Non-treated HeLa and UV treated HeLa cells were subjected to SDS PAGE followed by western blot with 81740-1-RR (Phospho-CHEK2 (Thr68) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin antibody as loading control. This data was developed using the same antibody clone with 81740-1-PBS in a different storage buffer formulation.