

For Research Use Only

Angiopoietin 1 Recombinant antibody

Catalog Number: 81990-4-RR



Basic Information

Catalog Number: 81990-4-RR	GenBank Accession Number: BC152411	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 284	CloneNo.: 240764D10
Source: Rabbit	UNIPROT ID: Q15389	Recommended Dilutions: WB 1:1000-1:4000
Isotype: IgG	Full Name: angiopoietin 1	
Immunogen Catalog Number: AG18829	Calculated MW: 498 aa, 58 kDa	
	Observed MW: 70 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : A549 cells, HeLa cells, U-87 MG cells
Species Specificity: human	

Background Information

Angiopoietin 1 is a 70-kDa secreted glycoprotein generated from vascular mural cells, pericytes, and certain other cells. Angiopoietin 1 participates in vascular differentiation through angiogenesis, which is the process of the growth and remodelling of existing vessels. Angiopoietin 1 is also involved in the maintenance and turnover of blood vessels in mature animals. Angiopoietin 1 binds to and activates the TEK/TIE2 receptor by inducing its dimerization and tyrosine phosphorylation, playing an important role in the regulation of angiogenesis. Overexpression of Angiopoietin 1 has been proven to occur in malignant glioblastoma, neuroblastoma, non-small cell lung cancer, and other tumors.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81990-4-RR (Angiopoietin 1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



1×10^6 HepG2 cells were intracellularly stained with 0.25 μ g Angiopoietin 1 Recombinant antibody (81990-4-RR, Clone:240764D10) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 μ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Bioluminescence resonance energy transfer (BRET) kinetic assays of 81990-4-RR against Human ANGPT1 were performed. The affinity constant is 6.37 nM.