

For Research Use Only

# Phospho-PERK/EIF2AK3 (Thr982) Recombinant antibody, PBS Only



Catalog Number: 82534-1-PBS

## Basic Information

Catalog Number:

82534-1-PBS

Size:

1mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC126354

GeneID (NCBI):

9451

UNIPROT ID:

Q9NZJ5

Full Name:

eukaryotic translation initiation  
factor 2-alpha kinase 3

Calculated MW:

1116 aa, 125 kDa

Observed MW:

180 kDa

Purification Method:

Protein A purification

CloneNo.:

4E16

## Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

Human

## Background Information

EIF2AK3 encodes the protein kinase RNA-like ER kinase (PERK), a key regulator of the unfolded protein response (UPR) in response to ER stress. Under ER stress conditions, activation of PERK is triggered by the dissociation of glucose-regulated protein (GRP) 78 (also known as BiP) from its luminal domain, followed by oligomerization and autophosphorylation. Phosphorylated PERK subsequently phosphorylates eukaryotic translation initiation factor 2 alpha (eif2  $\alpha$ ), to attenuate global protein translation and reduce incoming ER protein load via upregulated ER chaperone expression. (PMID: 35922637, PMID: 32029570)

## Storage

Storage:

Store at -80°C.

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Non-treated HEK-293 cells and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 82534-1-RR (Phospho-PERK/EIF2AK3 (Thr982) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with beta actin antibody (66009-1-Ig) as loading control. This data was developed using the same antibody clone with 82534-1-PBS in a different storage buffer formulation.



Non-treated NIH/3T3 cells and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 82534-1-RR (Phospho-PERK/EIF2AK3 (Thr982) antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82534-1-PBS in a different storage buffer formulation.