## For Research Use Only

## Cyclin D1 Recombinant antibody

Catalog Number:82681-1-RR



**Basic Information** 

Catalog Number: GenBank Accession Number: 82681-1-RR BC00076

 Size:
 GeneID (NCBI):

 1000 μg/ml
 595

Source: UNIPROT ID:
Rabbit P24385
Isotype: Full Name:
IgG cyclin D1

Immunogen Catalog Number: Calculated MW: AG25384 295 aa, 34 kDa

Observed MW: 34 kDa

**Applications** 

**Tested Applications:** 

IHC, WB, ELISA
Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: HepG2 cells, A431 cells, HCT 116 cells, SW 1990 cells, NIH/3T3 cells, PC-12 cells

**Purification Method:** 

CloneNo.:

6B10

Protein A purification

Recommended Dilutions: WB 1:5000-1:50000

IHC 1:500-1:2000

IHC: human oesophagus cancer tissue,

**Background Information** 

CCND1 (cyclin D1), also known as PRAD1 or BCL1, belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. CCND1 forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. The CCND1 gene, located on 11q13 has been reported to be overexpressed in mantle cell lymphoma (MCL) due to the chromosomal translocation. CCND1 has been shown to interact with tumor suppressor protein Rb and the expression of this gene is regulated positively by Rb. Over-expression of CCND1 is known to correlate with the early onset of cancer and risk of tumor progression and metastasis.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82681-1-RR (Cyclin D1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 82681-1-RR (Cyclin D1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).