For Research Use Only

YY1 Recombinant antibody

Catalog Number:82712-3-RR



Basic Information

Catalog Number:

GenBank Accession Number: BC037308

Purification Method: Protein A purification

82712-3-RR

GeneID (NCBI):

Size:

CloneNo.:

400 μg/ml

7528 **UNIPROT ID:** P25490

Source: Rabbit

Full Name:

Recommended Dilutions: WB 1:1000-1:8000

Isotype:

YY1 transcription factor

IF 1:200-1:800

Immunogen Catalog Number: AG17792

Calculated MW: 414 aa, 45 kDa

Observed MW:

65 kDa

Applications

Tested Applications:

Positive Controls:

IF/ICC, WB, ELISA

WB: HeLa cells,

Species Specificity: Human, mouse

IF: HepG2 cells,

Background Information

YY1, also named as DELTA, INO 80S and NF-E1, contains four C2H2-type zinc fingers and belongs to the YY transcription factor family. YY1 is a multifunctional transcription factor that exhibits positive and negative control on a large number of cellular and viral genes by binding to sites overlapping the transcription start site. YY1 may direct histone deacetylases and histone acetyltransferases to a promoter in order to activate or repress the promoter, thus implicating histone modification in the YY1. The open reading frame of the human YY1 cDNA encodes a protein of 414 amino acids with a predicted molecular weight of 44 kDa. However, YY1 migrates on SDS gels as a 65-68 kDa protein, probably due to the structure of the protein. It is a ubiquitously expressed transcription factor with fundamental roles in embryogenesis, differentiation, replication and proliferation.

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82712-3-RR (YY1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using YY1 antibody (82712-3-RR, Clone: 1B5) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).