For Research Use Only

## BST2 Recombinant antibody, PBS Only

Catalog Number:82733-7-PBS

Featured Product



Basic Information	Catalog Number: 82733-7-PBS	GenBank Accession Number: BC033873	Purification Method: Protein A purification
	Size: 1 mg/ml	GenelD (NCBI): 684	CloneNo.: 7J24
	Source: Rabbit	UNIPROT ID: Q10589	
	sotype:Full Name:gGbone marrow stromal cell antigen 2		n 2
	Immunogen Catalog Number: AG4430	Calculated MW: 180 aa, 20 kDa	
		Observed MW: 30-36 kDa	
Applications	Tested Applications: WB, IF/ICC, FC (Intra), Indirect ELISA		
	Species Specificity: human		
Background Information	BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. BST2 may play a role in B-cell activation in rheumatoid arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane proteins, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flavoviruses whilst IFITMs disrupt influenza and flavivirus infection. Observed MW of BST2 is 30-36 kDa (PMID: 19196977; 21237475).		
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS Only		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 82733-7-RR (BST2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82733-7-PBS in a different storage buffer formulation.



WB result of BST2 antibody (82733-7-RR; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-BST2 transfected HeLa cells. This with 82733-7-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepC2 cells using BST2 antibody (82733-7-RR, Clone: 7124) at dilution of 1:125 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgC(H+L) (SA00013-2). This data was developed using the same antibody clone with 82733-7-PBS in a different storage buffer formulation formulation.

1x10^6 HepG2 cells were intracellularly stained with 0.25 ug BST2 Recombinant antibody (82733-7-RR, Clone:7J24) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgC(H+L) (SA00013-2) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 82733-7-PBS in a different storage buffer formulation.

formulation.



Biolayer interferometry (BLI) kinetic assays of 82733-7-RR against Human BST2 were performed. The affinity constant is below 1 pM.