## For Research Use Only

## spike protein Recombinant antibody

Catalog Number:82734-4-RR



**Purification Method:** 

CloneNo.:

4118

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000

**Basic Information** 

Catalog Number:

82734-4-RR NC\_045512 GeneID (NCBI):

250 µg/ml 43740568 Source: Full Name:

Rabbit SARS-CoV-2 Spike Protein

Calculated MW: Isotype: 141 kDa

Immunogen Catalog Number:

AG30684

Size:

Positive Controls:

WB: Recombinant protein,

**Applications** 

**Tested Applications:** WB, ELISA

Species Specificity:

Virus

## **Background Information**

Coronaviruses (CoVs) infect human and animals and cause varieties of diseases, including respiratory, enteric, renal, and neurological diseases. CoV uses its spike protein to recognize ACE2 as its receptors and mediate membrane fusion and virus entry into host cells(PMID: 32221306). Each monomer of trimeric S protein is about 180 kDa, and contains two subunits, S1 and S2,S1 recognizes and binds to host receptors, and subsequent conformational  $changes in S2\ facilitate\ fusion\ between\ the\ viral\ envelope\ and\ the\ host\ cell\ membrane\ (PMID:\ 19198616).\ Although\ membrane\ (PMID:\ 19198616).$ the amino acid sequences of the S-glycoprotein were found to be different between the various HCoV, the structures showed high similarity, but the best 3D structural overlap shared by SARS-CoV and SARS-CoV-2, consistent with the shared ACE2 predicted receptor (PMID: 32522207). The spike protein of CoVs can be a target for vaccine and therapeutic development (PMID: 19198616). 28901-1-AP is specific for spike protein of SARS-COV-2, that antigen region is 428-506aa.

GenBank Accession Number:

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Recombinant protein were subjected to SDS PAGE followed by western blot with 82734-4-RR (spike protein antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.