For Research Use Only

SIRT1 Recombinant antibody, PBS Only

Catalog Number:82790-4-PBS Featured Product



Basic Information

Catalog Number: 82790-4-PBS

Concentration:

BC012499 GeneID (NCBI): 23411 **UNIPROT ID:**

> Q96EB6 Full Name:

GenBank Accession Number:

Protein A purification CloneNo.: 2F11

Purification Method:

1mg/ml Source: Rabbit Isotype:

sirtuin (silent mating type

information regulation 2 homolog) 1 Immunogen Catalog Number: (S. cerevisiae)

> Calculated MW: 747 aa, 82 kDa Observed MW: 110-130 kDa

Applications

Tested Applications:

WB, IF/ICC, IP, Indirect ELISA

Species Specificity:

human

AG3808

Background Information

SIRT1, also named as SIR2L1, contains a deacetylase sirtuin-type domain and belongs to the sirtuin family. The posttranslation modified SIRT1 is a 110-130 kDa protein, which contains one deacetylase sirtuin-type domain. The 75-80 kDa SirT1 fragment was detected to lack the carboxy-terminus (PMID:21305533). SirT1 exists a 57-61 kDa isoform. SIRT1 may be found in nucleolus, nuclear euchromatin, heterochromatin, and inner membrane. It can shuttles between the nucleus and cytoplasm. SIRT1 regulates processes such as apoptosis and muscle differentiation by deacetylating key proteins. SIRT1 in particular initiates several signaling events relevant to cardioprotection, including activation of endothelial nitric oxide synthase, INS receptor signaling, and autophagy. In addition, SIRT1 activation elicits resistance to oxidative stress via the regulation of transcription factors and co-activators such as FOXO, Hif-2a, and NF-kB. SIRT1 regulates the p53-dependent DNA damage response pathway by binding to and deacetylating p53, specifically at Lysine 382.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS only, pH7.3


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82790-4-RR (SIRT1 antibody) at dilution of 1:4700 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SIRT1 antibody (82790-4-RR, Clone: 2F11) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004). This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.




WB result of SIRT1 antibody (82790-4-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SIRT1 transfected HeLa cells. This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SIRT1 antibody (82790-4-RR, Clone: 2F11) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.



IP result of anti-SIRT1 (IP:82790-4-RR, 4ug; Detection:82790-4-RR 1:1000) with HEK-293 cells lysate 1360 ug. This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 82790-4-RR against Human SIRT1 were performed. The affinity constant is 4.90 nM.