For Research Use Only

MYB/c-Myb Recombinant antibody, PBS Only




Catalog Number:82804-2-PBS


Basic Information	Catalog Number: 82804-2-PBS	GenBank Accession Number: BC064955	Purification Method: Protein A purification
	Size: 1mg/ml	Genel D (NCBI): 4602	CloneNo.: 230179C11
	Source: Rabbit	UNIPROT ID: P10242	
	lsotype: lgG Immunogen Catalog Number: AG12021	Full Name: v-myb myeloblastosis viral oncogene homolog (avian) Calculated MW: 640 aa, 72 kDa	
		Observed MW: 40-50 kDa, 72-80 kDa	
Applications	Tested Applications: Indirect ELISA, WB Species Specificity: human		
Background Information	The MYB protein is a transcription factor that plays a key role in regulating the proliferation/differentiation of progenitor cells in bone marrow, colonic crypts, and neurogenic niches [PMID:18574464]. It is involved in the regulation of epithelial-to-mesenchymal transition (EMT) and invasion in neuroblastoma, colon carcinoma, and embryonic kidney cells through the upregulation of the transcription repressor Slug [PMID:20622260]. It also a DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3', and plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells. MYB exists some isoforms with MV 85, 75, 72, 68, 63 and 40 kDa.		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	cks. Upon receipt, store it immediatel	y at −80°C

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 82804-2-RR (MYB antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82804-2-PBS in a different storage buffer formulation. 1x10^6 MCF-7 cells were intracellularly stained with 0.25 ug MYB/c-Myb Recombinant antibody (82804-2-RR, Clone:230179C11) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 82804-2-PBS in a different storage buffer formulation.