

EDNRB Recombinant antibody

Catalog Number: 82818-2-RR

Basic Information

Catalog Number:

82818-2-RR

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_000115

GeneID (NCBI):

1910

UNIPROT ID:

P24530

Full Name:

endothelin receptor type B

Calculated MW:

50 kDa

Observed MW:

60 kDa

Purification Method:

Protein A purification

CloneNo.:

102

Recommended Dilutions:

WB 1:1000-1:5000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB : HEK-293 cells, mouse brain tissue, mouse cerebellum tissue, rat brain tissue

Background Information

EDNRB (endothelin receptor type B) is a G-protein coupled heptahelical endothelin-1 (EDN1) receptor. EDN1, a known mediator of human myometrial contraction in vitro, belongs to a family of three 21-amino acid isopeptides (PMID:18497345). EDNRB is a 442-residue protein expressed post-natally in many tissues, particularly in the myenteric plexus, mucosal layer, ganglia, and blood vessels of the submucosa of the colon (PMID:11434563). Moreover, Both the EDNRA and EDNRB are expressed in vessels and can be bound by EDN1, but while EDNRA mediates a constrictive effect, EDNRB is involved in vasorelaxation and functions as a clearance receptor that scavenges circulating EDN1 (PMID:30171849).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82818-2-RR (EDNRB antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



1×10^6 Raji cells were intracellularly stained with 0.25 μ g EDNRB Recombinant antibody (82818-2-RR, Clone:1O2) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 μ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).