## For Research Use Only

## Mono-Methyl-Histone H3 (Lys9) Recombinant antibody

Antibodies | ELISA kits | Proteins www.ptglab.com


Catalog Number:82821-6-RR


| Basic Information      | Catalog Number:<br>82821-6-RR   | GenBank Accession Number:<br>BC066245                          | Purification Method:<br>Protein A purification  |  |
|------------------------|---|--|---|--|
|                        | Size:<br>1000 µ g/ml  | GenelD (NCBI):<br>8350   | CloneNo.:<br>3C23   |  |
|                        | Source:<br>Rabbit   | UNIPROT ID:<br>P68431  | Recommended Dilutions:<br>WB 1:5000-1:50000   |  |
|                        | Isotype:<br>IgG   | Full Name:<br>histone cluster 1, H3a<br>Observed MW:<br>15 kDa |   |  |
|                        |   |  |   |  |
| Applications           | Tested Applications:<br>WB, ELISA, Dot Blot   |  | Positive Controls:<br>WB : HeLa cells, HEK-293 cells, K-562 cells, THP-1 cells,<br>HSC-T6 cells |  |
|                        | Species Specificity:<br>Human, rat  |  |   |  |
| Background Information | Histones, including H1/H5 (linker histones), H2, H3, and H4 (core histones), are nucleic proteins which interact with<br>DNA to form the nucleosomes and play important roles in gene regulation and DNA replication. Histone proteins are<br>highly post-translationally modified while Histone H3 is the most extensively modified. Methylation of Histone H3<br>at lysine 9 is linked to transcriptional repression. |  |   |  |
| Storage                | Storage:<br>Store at -20°C. Stable for one<br>Storage Buffer:<br>PBS with 0.02% sodium azide<br>Aliquoting is unnecessary for   | e and 50% glycerol pH 7.3.                                     |   |  |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Dot blot analysis was used to confirm the specificity of 82821-6-RR Mono-Methyl-Histone H3 (Lys9) antibody. Peptides were spotted onto NC and probed with antibody at 1  $\mu$ g/ml. The amount of peptide (ug/mL) spotted is indicated next to each row.

Various lysates were subjected to SDS PAGE followed by western blot with 82821-6-RR (Mono-Methyl-Histone H3 (Lys9) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.