For Research Use Only

## RBP4 Recombinant antibody, PBS Only

Catalog Number:82934-1-PBS

Featured Product




Basic Information	Catalog Number: 82934-1-PBS	GenBank Accession Number: BC020633	Purification Method: Protein A purification
	Concentration: 1mg/ml	GenelD (NCBI): 5950	CloneNo.: 6013
	Source: Rabbit	UNIPROT ID: P02753	
	Isotype: IgG	Full Name: retinol binding protein 4, plasma	
	Immunogen Catalog Number: AG2448	Calculated MW: 201 aa, 23 kDa	
		Observed MW: 23 kDa	
Applications	Tested Applications: WB, FC (Intra), Indirect ELISA		
	Species Specificity: human		
Background Information	RBP4 (retinol-binding protein 4) is a carrier protein that transports vitamin A (retinol) from the liver to the peripheral tissues. Synthesized primarily by hepatocytes and adipocytes as a 21 kDa non-glycosylated protein, RBP4 is secreted into the circulation as a retinol-RBP4 complex. In plasma the RBP4-retinol complex is bound to transthyretin (TRR), which prevents prevent kidney filtration. Two truncated forms of RBP4, RBP4-L (truncated at Leu-183) and RBP4-LL (truncated at Leu-182 and Leu-183), exist by proteolytic process. RBP4-L and RBP4-LL, which do not bind TTR, are normally excreted into the urine but accumulate in the serum during renal failure. Urinary RBP4 has been reported as marker for glomerular disease. RBP4 could be used to assess the risk of INS resistance, type 2 diabetes, obesity, and cardiovascular disease. (18752671, 16034410)		
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS only, pH7.3		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



50BD→ 50BD→ 40BD→ 30BD→ 20BD→ 20BD→ 1400 1400 HepG2


Various lysates were subjected to SDS PAGE followed by western blot with 82934-1-RR (RBP4 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82934-1-PBS in a different storage buffer formulation.





٠

1x10<sup>^6</sup> HepG2 cells were intracellularly stained with 0.25 ug RBP4 Recombinant antibody (82934-1-RR, Clone:6013) and Coralite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 82934-1-PBS in a



Biolayer interferometry (BLI) kinetic assays of 82934-1-RR against Human RBP4 were performed. The affinity constant is below 1 pM.