For Research Use Only

## uPAR, PLAUR Recombinant antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

230148A1

Catalog Number:82953-1-PBS

**Basic Information** 

Catalog Number: 82953-1-PBS

Size: 1 mg/ml GenBank Accession Number:

BC002788

GeneID (NCBI):

5329

Source: UNIPROT ID:
Rabbit Q03405
Isotype: Full Name:
IgG PLAUR

Immunogen Catalog Number: Calculated MW:

AG28056 37 kDa

**Applications** 

Tested Applications: Indirect ELISA,FC Species Specificity:

Human

**Background Information** 


uPAR is a 45-65 kDa, highly glycosylated, GPI-anchored membrane protein. In addition to the membrane-anchored form, uPAR is released from the plasma membrane by cleavage of the GPI anchor and can be found as a soluble form (suPAR). uPAR contains three homologous domains (D1-D3) of which the N-terminal one (D1) represents the uPA-binding domain. After binding to uPAR, uPA cleaves plasminogen, generating the active protease plasmin which is involved in a wide variety of physiologic and pathologic processes. In addition to regulating proteolysis, uPAR has important function in cell adhesion, migration and proliferation. Studies reveal that uPAR expression is elevated during inflammation and tissue remodeling and in many human cancers, in which it frequently indicates poor prognosis. (PMID 20027185; 12461559)

Storage

Storage:

Store at -80°C. Storage Buffer: PBS Only

## Selected Validation Data



1x10^6 A431 cells were intracellularly stained with 0.25 ug Anti-Human uPAR, PLAUR (82953-1-RR, Clone: 230148A1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:166 (red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:166 (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same