For Research Use Only

IL-12/IL-23 p40 Recombinant antibody

Catalog Number:83008-7-RR



Basic Information

Catalog Number:

GenBank Accession Number: BC067498

Purification Method:

83008-7-RR

Protein A purfication

Size:

GeneID (NCBI):

CloneNo.: 241600B3

1000 µg/ml

UNIPROT ID:

Recommended Dilutions:

Source: Rabbit

P29460

IF/ICC 1:125-1:500

Isotype:

Full Name:

interleukin 12B (natural killer cell

stimulatory factor 2, cytotoxic

lymphocyte maturation factor 2, p40)

Calculated MW:

328 aa, 37 kDa

Applications

Tested Applications:

IF/ICC, ELISA

Species Specificity:

human

Positive Controls:

IF/ICC: MOLT-4 cells,

Background Information

 $IL-12 \ and \ IL-23 \ are \ heterodimeric \ cytokines \ that \ share \ a \ common \ p40 \ subunit \ (PMID: 11114383). \ IL-12 \ is \ composed$ of the IL-12 p40 subunit linked to the IL-12 p35 subunit, and the heterodimer signals through the IL-12 receptor (IL-12R), which comprises the IL-12R β 1 and IL-12R β 2 subunits. IL-23 is composed of the IL-23 p19 subunit and the IL-12 p40 (IL-12/23p40) subunit, which signals through IL-23R and IL-12R β 1 (PMID: 11114383; 26121196). IL-12/IL-23 p40 also exists as a monomer and as a homodimer which can act as a potent IL-12 antagonist (PMID: 8958912; 18783467). IL-12/IL-23 p40 is produced by antigen-presenting cells, such as dendritic cells (DCs), monocytes, macrophages, neutrophils and, to a lesser extent, B cells (PMID: 20476918).

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed MOLT-4 cells using IL12B antibody (83008-7-RR, Clone: 241600B3) at dilution of 1:250 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).