For Research Use Only

VEGF Receptor 2 Recombinant antibody



Catalog Number:83049-4-RR

Basic Information

Catalog Number:

GenBank Accession Number: NM_002253

Purification Method:

83049-4-RR

Protein A purification

Size:

GeneID (NCBI):

CloneNo.:

1000 µg/ml Source:

230238D7

Rabbit

UNIPROT ID: P35968-1

Recommended Dilutions: WB 1:500-1:2000

Isotype:

Full Name: kinase insert domain receptor (a type

IHC 1:200-1:800

III receptor tyrosine kinase)

Calculated MW:

152kd

Observed MW:

150 kDa, 230 kDa

Applications

Tested Applications:

Positive Controls:

WB, IHC, ELISA

Species Specificity:

WB: human placenta tissue, IHC: human placenta tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

KDR, also named as VEGFR-2, FLK1 and CD309, is a receptor for VEGF or VEGFC. KDR which belongs to the protein kinase superfamily, has a tyrosine-protein kinase activity. The VEGF-kinase ligand/receptor signaling system plays a key role in vascular development and regulation of vascular permeability. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions. KDR functions as the main mediator of VEGF-induced endothelial proliferation, survival, migration, tubular morphogenesis and sprouting. Mutations of this gene are implicated in infantile capillary hemangiomas.

Storage


Storage:


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data








Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 83049-4-RR (KDR antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



human placenta tissue was subjected to SDS PAGE followed by western blot with 83049-4-RR (KDR antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLL) kinetic assays of 83049-4-RR against Human KDR were performed. The affinity constant is 4.71 nM.