

For Research Use Only

Integrin beta 3 Recombinant antibody



Catalog Number: 83053-6-RR

Basic Information

Catalog Number: 83053-6-RR	GenBank Accession Number: BC127667	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 3690	CloneNo.: 230292A10
Source: Rabbit	UNIPROT ID: P05106	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: integrin, beta 3 (platelet glycoprotein IIIa, antigen CD61)	
	Calculated MW: 87 kDa	
	Observed MW: 100-110 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : human peripheral blood platelets,
Species Specificity: Human	

Background Information

Integrin beta-3, also named as CD61 and GPIIIa, is a receptor for cytotactin, fibronectin, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin, vitronectin and von Willebrand factor. Integrin beta-3 recognize the sequence R-G-D in a wide array of ligands and the sequence H-H-L-G-G-A-K-Q-A-G-D-V in fibrinogen gamma chain. Following activation integrin beta-3 brings about platelet/platelet interaction through binding of soluble fibrinogen which leads to rapid platelet aggregation which physically plugs ruptured endothelial surface. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions. Human integrin beta-3 has a calculated molecular weight of 87 kDa. As a result of glycosylation, the apparent molecular mass of integrin beta-3 is approximately 90-110 kDa.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 83053-6-RR (Integrin beta-3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.