## For Research Use Only

## NEDD4 Recombinant antibody

Catalog Number:83112-4-RR



**Basic Information** 

Catalog Number: GenBank Accession Number: 83112-4-RR BC152452

 Size:
 GeneID (NCBI):

 1000 μg/ml
 4734

 Source:
 UNIPROT ID:

 Rabbit
 P46934

IgG neural precursor cell expressed,
Immunogen Catalog Number: developmentally down-regulated 4

AG16291 Calculated MW:

1319 aa, 149 kDa

Observed MW: 115 kDa

Full Name:

**Applications** 

Tested Applications: WB, IF/ICC, ELISA

Isotype:

Species Specificity: Human, Mouse, Rat Positive Controls:

WB: C6 cells, A549 cells, Jurkat cells, NIH/3T3 cells

**Purification Method:** 

CloneNo.:

230478D8

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IF 1:125-1:500

IF: U2OS cells,

## **Background Information**

NEDD4 and similar proteins discovered subsequently became a family of HECT ligases, comprising 9 human proteins including NEDD4, NEDD4-2 (NEDD4L), ITCH, SMURF1, SMURF2, WWP1, WWP2, NEDL1 and NEDL2. NEDD4 is a highly evolutionarily conserved protein from yeast to man, and was initially cloned as a highly expressed gene in the early embryonic brain. NEDD4 is frequently overexpressed in many different types of cancer, and decreased levels of NEDD4 can also be associated with cancer. It can be a potential therapeutic target for the treatment of human cancer.(PMID: 25527121). The human NEDD4 gene is located on chromosome 15q21.3 and comprises 30 exons (HGNC:7727) shown to encode a ~120 KDa protein. Otherwise there is a 75 kDa isoform in addition to full length NEDD4 (PMID: 24907641).

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 83112-4-RR (NEDD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 83112-4-RR (NEDD4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunofluorescent analysis of (-20°C Ethanol) fixed U2OS cells using NEDD4 antibody (83112-4-RR, Clone: 230478D8 ) at dilution of 1:250 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).