For Research Use Only

## SAR1A Recombinant antibody

Catalog Number:83168-6-RR




Size: 1000 µg/ml	GenelD (NCBI):	Protein A purfication	
	56681	CloneNo.: 230471A11	
Source: Rabbit	UNIPROT ID: Q9NR31	Recommended Dilutions: WB 1:5000-1:50000	
Isotype: IgG	Full Name: SAR1 homolog A (S. cerevisiae) Calculated MW: 22 kDa		
			Observed MW: 23-25 kDa
	Tested Applications: WB_EC (Intra)_ELISA		Positive Controls: WB : HEK-293T cells, NIH/3T3 cells, HeLa cells, Jurkat cells, mouse liver tissue
Species Specificity: human, mouse			
involved in transport from the localization at discrete locati	e endoplasmic reticulum to the Golgi appa ons on the ER membrane perhaps by preve	aratus. SAR1A is required to maintain SEC16/ enting its dissociation. SAR1A-GTP-	
Storage Buffer:			
	Isotype: IgG Tested Applications: WB, FC (Intra), ELISA Species Specificity: human, mouse SAR1A, also named as SAR1, involved in transport from the localization at discrete locati dependent assembly of SEC1 exit sites (ERES). Storage: Storage Buffer: PBS with 0.02% sodium azide	Isotype: Full Name:   IgG SAR1 homolog A (S. cerevisiae)   Calculated MW: 22 kDa   Observed MW: 23-25 kDa   Tested Applications: Positive Q   WB, FC (Intra), ELISA WB : HEK-   Species Specificity: cells, mothuman, mouse   SAR1A, also named as SAR1, SARA and SARA1, belongs to the small GT involved in transport from the endoplasmic reticulum to the Golgi apparlocalization at discrete locations on the ER membrane perhaps by previdependent assembly of SEC 16A on the ER membrane forms an organizate exit sites (ERES).   Storage: Storage:   Storage: Storage for one year after shipment.	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data

temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 83168-6-RR (SAR1A antibody) at dilution of 1:10000 incubated at room

1x10^6 A549 cells were intracellularly stained with 0.25 ug SAR1A Recombinant antibody (83168-6-RR, Clone:230471A11) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).

10<sup>3</sup> 10<sup>4</sup>

83168-6-RR(SAR1A),APC-H

10<sup>5</sup> 10<sup>6</sup> 10

100-


80

60

40

20

Relative Cell Number



1x10^6 A431 cells were intracellularly stained with 0.25 ug SAR1A Recombinant antibody (83168-6-RR, Clone:230471A11) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).