For Research Use Only

SNAP25 Recombinant antibody

Catalog Number:83259-1-RR



Basic Information

Catalog Number:

GenBank Accession Number: BC010647

Purification Method:

83259-1-RR

Protein A purification

Size: 1000 µg/ml GeneID (NCBI): 6616

CloneNo.: 240069A1

Source:

UNIPROT ID: P60880

Recommended Dilutions:

Rabbit Isotype:

Full Name:

IF 1:125-1:500

AG6695

synaptosomal-associated protein,

25kDa Immunogen Catalog Number:

Calculated MW:

23 kDa

Applications

Tested Applications: IF/ICC, FC, ELISA

IF: PC-12 cells,

Positive Controls:

Species Specificity:

Human, Rat

Background Information

The synaptosomal associated protein of 25 kD (SNAP-25) was first identified as a major synaptic protein by Wilson and colleagues. The protein interacts with syntaxin and synaptobrevin through its N-terminal and C-terminal helical domains. Its palmitoylation domain is located in the middle of the molecule that contains four cysteine residues. Mutation of the cysteines abolishes palmitoylation and membrane binding. Several elegant studies using synaptosome preparations and permeabilized PC12 cells have suggested that SNAP-25 may act in the late postdocking steps of exocytosis. By limited proteolysis and in vitro binding assay, it is proposed that the two helix domains act independently and contribute equally to form the SNARE complex with syntaxin and synaptobrevin. It seems that a major regulatory element is located in the C-terminus of SNAP-25. Removing a 9 amino acid sequence of SNAP-25 inhibited neurosecretion in chromaffin cells. In addition, it has been shown that inhibition of neurosecretion by AX type E can be rescued by a SNAP-25 C-terminal peptide, probably by initiating the formation of a fusion competent SNARE complex.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



1x10^6 PC-12 cells were intracellularly stained with 0.25 ug SNAP25 Recombinant antibody (83259-1-RR, Clone:240069A1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (4% PFA) fixed PC-12 cells using SNAP antibody (83259-1-RR, Clone: 240069A1) at dilution of 1:250 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).