

For Research Use Only

# Nucleolin/C23 Recombinant antibody



Catalog Number: 83380-1-RR

## Basic Information

**Catalog Number:**

83380-1-RR

**Size:**

800 µg/ml

**Source:**

Rabbit

**Isotype:**

IgG

**Immunogen Catalog Number:**

AG0859

**GenBank Accession Number:**

BC006494

**GeneID (NCBI):**

4691

**UNIPROT ID:**

P19338

**Full Name:**

nucleolin

**Calculated MW:**

76 kDa

**Observed MW:**

100-110 kDa

**Purification Method:**

Protein A purification

**CloneNo.:**

240029F8

**Recommended Dilutions:**

WB 1:5000-1:50000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IF 1:125-1:500

## Applications

**Tested Applications:**

WB, IP, IF/ICC, ELISA

**Species Specificity:**

Human

**Positive Controls:**

WB : HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, MCF-7 cells

IP : HeLa cells,

IF : HepG2 cells,

## Background Information

Nucleolin, also known as C23, involved in the control of transcription of ribosomal RNA (rRNA) genes by RNA polymerase I, in nucleocytoplasmic transportation of ribosomal components, and in ribosome maturation and assembly. It associated with intranucleolar chromatin and pre-ribosomal particles, and induced chromatin decondensation by binding to histone H1. Also it has a role in the process of transcriptional elongation. Whilst mammalian nucleolin has a predicted molecular mass of approximately 77 kDa, the apparent molecular mass is between 100 and 110 kDa, and has been attributed to the amino acid composition of the N-terminal domain, which is highly phosphorylated. (PMID: 15925566)

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83380-1-RR (Nucleolin/C23 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



IP result of anti-Nucleolin/C23 (IP:83380-1-RR, 4ug; Detection:83380-1-RR 1:5000) with HeLa cells lysate 1085 ug.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using Nucleolin/C23 antibody (83380-1-RR, Clone: 240029F8 ) at dilution of 1:250 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Biolayer interferometry (BLI) kinetic assays of 83380-1-RR against Human Nucleolin/C23 were performed. The affinity constant is 2.47 nM.