

For Research Use Only

SIAH1 Recombinant antibody

Catalog Number: 83389-4-RR



Basic Information

Catalog Number: 83389-4-RR	GenBank Accession Number: BC035562	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 6477	CloneNo.: 240377B4
Source: Rabbit	UNIPROT ID: Q8IUQ4	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG	Full Name: seven in absentia homolog 1 (Drosophila)	
Immunogen Catalog Number: AG34618	Calculated MW: 282 aa, 31 kDa	
	Observed MW: 30-34 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB: A549 cells, mouse kidney tissue, HepG2 cells, HEK-293T cells, SH-SY5Y cells, Jurkat cells, COLO 320 cells
Species Specificity: Human, mouse	

Background Information

SIAH1, also named as HUMSIAH, Siah-1a and Siah-1S(21kd), belongs to the SINA (Seven in absentia) family. SIAH1 is known to cause indirect degradation of beta-catenin through formation of a complex with Siah-interacting protein (SIP), Skp1 and Ebi. SIAH1 can form heterodimer or homodimer such as Siah-1^{*}Siah-1(70 or 62kd), Siah-1^{*}Siah-1S(42kd) and Siah-1S^{*}Siah-1S(51-56kd). Importantly, results from in vitro soft agar assay demonstrated that Siah-1S displays a promotion effect on cells tumorigenicity. (PMID:17420721)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83389-4-RR (SIAH1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



mouse kidney tissue were subjected to SDS PAGE followed by western blot with 83389-4-RR (SIAH1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.