## For Research Use Only

## PMS1 Recombinant antibody

Catalog Number:83456-6-RR



**Basic Information** 

Catalog Number: GenBank Accession Number: 83456-6-RR BC096331

 Size:
 GeneID (NCBI):

 1000 μ g/ml
 5378

 Source:
 UNIPROT ID:

 Rabbit
 P54277

 Isotype:
 Full Name:

IgG PMS1 postmeiotic segregation increased 1 (S. cerevisiae)

AG28132 Observed MW: 106 kDa

**Applications** 

Tested Applications: WB, IF/ICC, FC (Intra), ELISA Species Specificity:

human

Purification Method: Protein A purfication

CloneNo.: 240408D9

Recommended Dilutions: WB 1:500-1:2000 IF/ICC 1:125-1:500

Positive Controls:

WB: HeLa cells, HepG2 cells, HEK-293T cells, K-562 cells, Jurkat cells, MOLT-4 cells, HL-60 cells

IF/ICC: Caco-2 cells, HeLa cells

## **Background Information**

PMS1 Homolog 1 is a member of the DNA mismatch repair enzymes (MMREs) which is to eliminate the mismatch of insertions and deletions as a consequence of DNA polymerase errors at DNA synthesis in eukaryotes. PMS1 and MLH1 could form a heterodimer Mutl.  $\beta$  that belongs to MMREs (PMID:25619773). In the MMR system, the MutS complex recognizes the DNA mispairs firstly, and then the Mlh1-Pms1 and other associated proteins such as PCNA, RFC, Exol were recruited, inducing Exonuclease 1 (Exo1)-dependent and -independent MMR pathways (PMID:24981171). Mutations in this gene cause hereditary nonpolyposis colorectal cancer (HNPCC), which is also known as Lynch syndrome. It was reported that PMS1 is a widely expressed protein and located in the nucleus. Alternative splicing allows for 4 isoforms of PMS1 protein.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 83456-6-RR (PMS1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed Caco-2 cells using PMS1 antibody (83456-6-RR, Clone: 240408D9) at dilution of 1:250 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug PMS1 Recombinant antibody (83456-6-RR, Clone:240408D9) and APC-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H-L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLI) kinetic assays of 83456-6-RR against Human PMS1 were performed. The affinity constant is 0.367 nM.