For Research Use Only

HSPA2 Recombinant antibody

Catalog Number:83466-1-RR

Featured Product



Basic Information

Catalog Number: 83466-1-RR

Size:

Source: Rabbit

1000 ug/ml

BC036107 GeneID (NCBI): 3306 UNIPROT ID: P54652

GenBank Accession Number:

Isotype: Full Name:
IgG heat shock 70kDa protein 2

Immunogen Catalog Number: Calculated MW:
AG3539 639 aa, 70 kDa
Observed MW:

70 kDa

Purification Method: Protein A purification

CloneNo.: 240421A2

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000

Applications

Tested Applications: WB, IHC, ELISA Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HeLa cells, A431 cells, A549 cells, HaCaT cells,

NIH/3T3 cells, human kidney tissue

IHC: mouse testis tissue,

Background Information

Human HSPA2 is a member of the HSPA (HSP70) family of heat-shock proteins, encoded by the gene originally described as testis-specific. Recently, it has been reported that HSPA2 can be also expressed in human somatic tissues in a cell-type specific manner. HSPA2 is a cytosol/nuclear protein able to translocate between cytoplasm and nucleus. At physiological temperature, the HSPA2 was localized primarily in cytoplasm whereas, during heat sokel, localization shifted to nucleus and nucleoli (PMID: 18452162). HSPA2 is involved in sperm maturity, function and fertility. Aberrant expression of HSPA2 in testes has been connected with male infertility. Recently, HSPA2 has attracted increased interest due to its possible involvement in carcinogenesis of non-testicular tissues. This antibody well recognized the endogenous HSPA2 protein in testis and multiple cell lines. (21373891)

Storage


Storage:

Store at -20°C. Stable for one year after shipment.
Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83466-1-RR (HSPA2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




WB result of HSPA2 antibody (83466-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HSPA2 transfected HeLa cells.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 83466-1-RR (HSPA2 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 83466-1-RR (HSPA2 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 83466-1-RR against Human HSPA2 were performed. The affinity constant is 1.86 nM.