

For Research Use Only

HMGB2 Recombinant antibody

Catalog Number: 83482-4-RR



Basic Information

Catalog Number: 83482-4-RR	GenBank Accession Number: BC001063	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 3148	CloneNo.: 240453C11
Source: Rabbit	UNIPROT ID: P26583	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: high-mobility group box 2	
Immunogen Catalog Number: AG6135	Calculated MW: 24 kDa	
	Observed MW: 24-28 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : LNCaP cells, HeLa cells, HEK-293 cells, MCF-7 cells, Jurkat cells, NIH/3T3 cells, HSC-T6 cells
Species Specificity: human, mouse, rat	

Background Information

High mobility group protein B2 (HMGB2) belongs to a family of highly conserved proteins that contain HMG box domains (11246022,14871457). All three family members (HMGB1, HMGB2, and HMGB3) contain two HMG box domains and a C-terminal acidic domain. HMGB1 is a widely expressed and highly abundant protein (14871457). HMGB2 is widely expressed during embryonic development, but it is restricted to lymphoid organs and testis in adult animals (11262228). HMGB3 is only expressed during embryogenesis (9598312). While expression varies, the biochemical properties of the different family members may be indistinguishable. The HMG box domains facilitate the binding of HMGB proteins to the minor groove of DNA, which results in local bending of the DNA double helix. HMGB proteins are recruited by and help facilitate the assembly of site-specific DNA binding proteins to their cognate binding sites in chromatin. For example, HMGB1 and HMGB2 facilitate the binding of Hox proteins, Oct proteins, p53, Rel proteins, and steroid hormone receptor proteins to their target gene promoters (11246022,14871457). Furthermore, HMGB2 interacts with RAG1 to facilitate RAG complex binding to the recombinant signal sequence (RSS) and stimulate DNA-bending and subsequent VDJ cleavage at antigen receptor genes (19317908,10490593). In addition to their functions in the nucleus, HMGB proteins play a significant role in extracellular signaling associated with inflammation. HMGB2 is secreted by myeloid cells and promotes proliferation and migration of endothelial cells by binding to the receptor for advanced glycation endproducts (RAGE) (19811285). Research studies have shown that HMGB2 overexpression in hepatocellular carcinoma is associated with poor prognosis and shorter survival time (20851854). The calculated molecular weight of HMGB2 is 24 kDa, and the post-modification of HMGB2 is about 33-35 kDa. (18218727)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83482-4-RR (HMGB2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1×10^6 HeLa cells were intracellularly stained with 0.25 μ g HMGB2 Recombinant antibody (83482-4-RR, Clone:240453C11) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 μ g Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biacore BLI kinetic assays of 83482-4-RR against Human HMGB2 were performed. The affinity constant is 16.4 nM.