For Research Use Only

## NCAM2 Recombinant antibody

Catalog Number:83494-2-RR



Basic Information	Catalog Number: 83494-2-RR	GenBank Accession Number: BC052946	Purification Method: Protein A purification
	Size: 1000 µg/ml	GenelD (NCBI): 4685	CloneNo.: 240398C5
	Source: Rabbit	UNIPROT ID: O 15394	Recommended Dilutions: WB 1:5000-1:50000
	lsotype: IgG	Full Name: neural cell adhesion molecule 2	
	Immunogen Catalog Number: AG4821	Calculated MW: 93 kDa	
		Observed MW: 90 kDa, 125 kDa	
Applications	Tested Applications: WB, ELISA Species Specificity: Human, mouse , rat	Positive	Controls:
		WB : mouse brain tissue, mouse cerebellum tissue, ra brain tissue	
Background Information	The mammalian NCAM family has two members, NCAM1 and NCAM2 (PMID: 9729879; 20439718). Neural cell adhesion molecule 2 (NCAM2, also known as NCAM21) is a single-pass type I membrane protein that is highly expressed in the brain, which may play important roles in selective fasciculation and zone-to-zone projection of the primary olfactory axons or in neurodegenerative diseases such as Alzheimer's disease (PMID: 26611261). NCAM2 has two isoforms: NCAM2.1 bearing transmembrane and cytoplasmic domains and the shorter NCAM2.2 isoform, which is GPI-anchored (PMID: 9334170; 9221781). The molecular weight of NCAM2 is around 125 kDa and 93 kDa (PMID: 9334170; 3171685).		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data

310 kDa-245 kDa


180 kDa

140 kDa-

100 kDa-

75 kDa-

60 kDa



Various lysates were subjected to SDS PAGE followed by western blot with 83494-2-RR (NCAM2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. Biolayer interferometry (BLI) kinetic assays of 83494-2-RR against Human NCAM2 were performed. The affinity constant is 0.882 nM.