

For Research Use Only

CKB Recombinant antibody

Catalog Number: 83522-5-RR



Basic Information

Catalog Number: 83522-5-RR	GenBank Accession Number: BC001190	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 1152	CloneNo.: 240418D2
Source: Rabbit	UNIPROT ID: P12277	Recommended Dilutions: WB 1:500-1:3000
Isotype: IgG	Full Name: creatine kinase, brain	
Immunogen Catalog Number: AG7285	Calculated MW: 43 kDa	
	Observed MW: 40-45 kDa	

Applications

Tested Applications: WB, FC (Intra), ELISA	Positive Controls: WB : HeLa cells, SH-SY5Y cells, Y79 cells, mouse brain tissue, rat brain tissue
Species Specificity: human, mouse, rat	

Background Information

CKBB, also named as B-CK and CKB, is a member of the ATP:guanido phosphotransferase protein family. It is a cytoplasmic enzyme involved in energy homeostasis. CKBB reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such as creatine phosphate. It acts as a homodimer in brain as well as in other tissues, and as a heterodimer with a similar muscle isozyme in heart. CK isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa. CK MB consists of a dimer of nonidentical chains. With MM being the major form in skeletal muscle and myocardium, MB existing in myocardium, and BB existing in many tissues, especially brain. This antibody can recognize both CKB and CKM due to the high homology.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83522-5-RR (CKB antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



1×10^6 HeLa cells were intracellularly stained with 0.25 μ g CKB Recombinant Antibody (83522-5-RR, Clone:240418D2) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 μ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).