For Research Use Only

## MED12L Recombinant antibody

Catalog Number:83540-1-RR




Basic Information	Catalog Number: 83540-1-RR	GenBank Accession Number: NM_053002	Purification Method: Protein A purfication	
	Size: 1000 ug/ml	GenelD (NCBI): 116931	CloneNo.: 240101C11	
	Source: Rabbit	UNIPROT ID: Q86YW9 Full Name: mediator complex subunit 12-like	Recommended Dilutions: IHC 1:500-1:2000 IF/ICC 1:150-1:600	
	lsotype: IgG			
	Immunogen Catalog Number: AG34636	Calculated MW: 240 kDa		
Applications	Tested Applications: IHC, IF/ICC, FC (Intra), ELISA			
	TE buffer pH 9.0; (*) Altern	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0		
Background Informati	family. MED12L may be a compor of nearly all RNA polymerase II-d from gene-specific regulatory pro to promoters by direct interaction preinitiation complex with RNA p	MED12L, also named as KIAA1635, TNRC11L, TRALP and TRALPUSH, belongs to the Mediator complex subunit 12 family. MED12L may be a component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. MED12L is a mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. MED12L is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors. This antibody can recognize all the isoform of MED12L This antibody has no cross-reaction to MED12.		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20	1 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using MED12L antibody (83540-1-RR, Clone: 240101C11) at dilution of 1:300 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 83540-1-RR (MED12L antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 U2OS cells were intracellularly stained with 0.25 ug MED12L Recombinant antibody (83540-1-RR, Clone:240101C11) and APC-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).