

For Research Use Only

TIGIT Recombinant antibody

Catalog Number: 83545-1-RR



Basic Information

Catalog Number:

83545-1-RR

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_173799.4

GeneID (NCBI):

201633

UNIPROT ID:

Q495A1-1

Full Name:

T cell immunoreceptor with Ig and ITIM domains

Calculated MW:

26 kDa

Observed MW:

44 kDa

Purification Method:

Protein A purification

CloneNo.:

240423D3

Recommended Dilutions:

WB 1:1000-1:6000

IF/ICC 1:125-1:500

Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:

Human, mouse, rat

Positive Controls:

WB : Jurkat cells, NK-92 cells, NIH/3T3 cells, RAW 264.7 cells, EL-4 cells, rat spleen tissue

IF/ICC : Jurkat cells,

Background Information

TIGIT (T-cell immunoreceptor with Ig and ITIM domains), also known as VSIG9 or VSTM3, is an immune receptor expressed on T cells, including Treg and memory subsets, as well as on NK cells (PMID: 19011627). It contains an immunoglobulin variable domain, a transmembrane domain and an immunoreceptor tyrosine-based inhibitory motif (ITIM). TIGIT binds to poliovirus receptor (PVR, also called CD155) with high affinity, and also to PVRL2 (CD112) with lower affinity. The interaction of TIGIT with PVR on dendritic cells increases the secretion of IL-10 and decreases the secretion of proinflammatory cytokine and suppresses T-cell activation by promoting the generation of mature immunoregulatory dendritic cells. TIGIT can inhibit NK cytotoxicity directly through its ITIM (PMID: 19815499).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83545-1-RR (TIGIT antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using TIGIT antibody (83545-1-RR, Clone: 240423D3) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).