

For Research Use Only

Chemerin Recombinant antibody

Catalog Number: 83550-4-RR



Basic Information

Catalog Number:

83550-4-RR

Size:

800 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_001347168.1

GeneID (NCBI):

71660

UNIPROT ID:

Q9DD06

Full Name:

retinoic acid receptor responder
(tazarotene induced) 2

Calculated MW:

18 kDa

Observed MW:

14-19 kDa

Purification Method:

Protein A purification

CloneNo.:

240355A12

Recommended Dilutions:

WB 1:2000-1:16000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

mouse

Positive Controls:

WB : mouse adipose tissue, mouse brown adipose tissue

Background Information

Chemerin, also named as TIG2 and RARRES2, is adipocyte-secreted protein (adipokine) that regulates adipogenesis, metabolism and inflammation through activation of the chemokine-like receptor 1 (CMKLR1). Its other ligands include G protein-coupled receptor 1 (GPR1) and chemokine receptor-like 2 (CCRL2). It positively regulates adipocyte differentiation, modulates the expression of adipocyte genes involved in lipid and glucose metabolism and might play a role in angiogenesis, a process essential for the expansion of white adipose tissue. RARRES2 have both pro- and anti-inflammatory properties depending on the modality of enzymatic cleavage by different classes of proteases. The MW of Chemerin is 14-19 kDa.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83550-4-RR (Chemerin antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



1×10^6 C57BL/6 mouse liver cells were intracellularly stained with 0.25 μ g Chemerin Recombinant antibody (83550-4-RR, Clone:240355A12) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 μ g Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83550-4-RR against Mouse Chemerin were performed. The affinity constant is 3.76 nM.