For Research Use Only

TCF7L1 Recombinant antibody

Catalog Number:83588-3-RR



Basic Information

Catalog Number: GenBank Accession Number: 83588-3-RR BC058894

 Size:
 GeneID (NCBI):

 1000 ug/ml
 83439

 Source:
 UNIPROT ID:

 Rabbit
 Q9HCS4

Isotype: Full Name:

IgG transcription factor 7-like 1 (T-cell specific, HMG-box)

Immunogen Catalog Number: specific, HMG-bo AG34253 Calculated MW: 63 kDa

> Observed MW: 63 kDa, 75 kDa

Applications

Tested Applications: WB, FC (Intra), ELISA

Species Specificity: human, mouse

Positive Controls:

WB: A431 cells, HeLa cells, HEK-293 cells, NCCIT cells,

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000

CloneNo.:

240544F1

NIH/3T3 cells

Background Information

TCF7L1 is one of TCF/LEF transcription factors that are mediators of the Wnt signaling pathway and are antagonized by the TGF-beta signaling pathway. It binds to DNA and acts as a repressor in the absence of CTNNB1, and as an activator in its presence. It's necessary for the terminal differentiation of epidermal cells, the formation of keratohyalin granules and the development of the barrier function of the epidermis

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83588-3-RR (TCF7L1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



1x10^6 A431 cells were intracellularly stained with 0.25 ug TCF7L1 Recombinant antibody (83588-3-RR, Clone:240544F1) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).