For Research Use Only

## Alpha-1-microglobulin Recombinant antibody



Catalog Number:83678-3-RR

| Basic Information      | Catalog Number:<br>83678-3-RR  | GenBank Accession Number:<br>NM_001633.4  | Purification Method:<br>Protein A purfication               |  |
|------------------------|--|---|---|--|
|                        | Size:<br>1000 ug/ml  | GenelD (NCBI):<br>259   | CloneNo.:<br>240751G7                                       |  |
|                        | Source:<br>Rabbit  | UNIPROT ID:<br>P02760   | Recommended Dilutions:<br>WB 1:5000-1:50000                 |  |
|                        | Isotype:<br>IgG  | Full Name:<br>alpha-1-microglobulin/bikunin<br>precursor<br>Calculated MW:<br>39kDa |   |  |
|                        |  |   |   |  |
|                        |  | Observed MW:<br>30 kDa  |   |  |
| Applications           | Tested Applications:<br>WB, FC (Intra), ELISA  |   | Positive Controls:<br>WB : human plasma, human urine sample |  |
|                        | Species Specificity:<br>human  | WD. Hum   |   |  |
| Background Information | Alpha 1 microglobulin, also known as HI30, is a 27-30 kDa glycoprotein, present in various body fluids. It belongs to the lipocalin superfamily of hydrophobic ligand-binding proteins forming an internal ligand-binding pocket. The protein acts as a mediator of bacterial adhesion to polymer surfaces and is involved in inhibiting renal lithogenesis. |   |   |  |
| Storage                | Storage:<br>Store at -20°C. Stable for one<br>Storage Buffer:<br>PBS with 0.02% sodium azide<br>Aliquoting is unnecessary for  | e and 50% glycerol pH 7.3.  |   |  |


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




human plasma were subjected to SDS PAGE followed by western blot with 83678-3-RR (Alpha-1-microglobulin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



human urine sample were subjected to SDS PAGE followed by western blot with 83678-3-RR (Alpha-1-microglobulin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1x10^6 MCF-7 cells were intracellularly stained with 0.25 ug Alpha 1 microglobulin Recombinant antibody (83678-3-RR, Clone:240751G7) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLl) kinetic assays of 83678-3-RR against Human Alpha 1 microglobulin were performed. The affinity constant is 6.59 nM.