

For Research Use Only

MOSC2 Recombinant antibody

Catalog Number:83705-2-RR



Basic Information

Catalog Number:

83705-2-RR

Size:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG20694

GenBank Accession Number:

BC011973

GeneID (NCBI):

54996

UNIPROT ID:

Q969Z3

Full Name:

MOCO sulphurase C-terminal domain containing 2

Calculated MW:

335 aa, 38 kDa

Observed MW:

35-38 kDa

Purification Method:

Protein A purification

CloneNo.:

240747G6

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:200-1:800

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse liver tissue, HUVEC cells, HEK-293 cells, mouse kidney tissue, rat liver tissue

IHC : human stomach cancer tissue,

Background Information

MOSC domain-containing protein 2 (also known as MOSC2), also known as MARC2, is a component of prodrug-converting system, reduces a multitude of N-hydroxylated prodrugs particularly amidoximes, leading to increased drug bioavailability. Also, MOSC2 may be involved in mitochondrial N(omega)-hydroxy-L-arginine (NOHA) reduction, regulating endogenous nitric oxide levels and biosynthesis. The reductase activity is regulated under adipogenic conditions, and down-regulation of the terminal component MOSC2 resulted in decreased lipid synthesis, suggesting a possible physiological role of this enzyme system and its component MOSC2 in lipogenesis(PMID: 22203676).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83705-2-RR (MOSC2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83705-2-RR (MOSC2 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83705-2-RR (MOSC2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 83705-2-RR against Human MOSC2 were performed. The affinity constant is 0.473 nM.