## For Research Use Only

## TNF-alpha Recombinant antibody

Catalog Number:83721-7-RR



**Basic Information** 

Catalog Number:

83721-7-RR Size:

1000 ug/ml

Source: Rabbit

Isotype:

GenBank Accession Number:

NM\_000594.4 GeneID (NCBI):

7124

ENSEMBL Gene ID: ENSG00000232810

UNIPROT ID: P01375 Full Name:

tumor necrosis factor (TNF

superfamily, member 2)

Calculated MW: 26 kDa Observed MW: 26 kDa Purification Method:

Protein A purfication

CloneNo.: 240668B4

Recommended Dilutions:

WB 1:2000-1:10000

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity:

human

Positive Controls:

WB: THP-1 cells, LPS treated THP-1 cells

## **Background Information**

TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, ins resistance, and cancer.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Untreated, LPS and Brefeldin A treated THP-1 cells were subjected to SDS PAGE followed by western blot with 83721-7-RR (TNF-alpha antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLL) kinetic assays of 83721-7-RR against Human TNF-alpha were performed. The affinity constant is 8.51 nM.