

For Research Use Only

Ager/Rage Recombinant antibody

Catalog Number: 83742-5-RR



Basic Information

Catalog Number: 83742-5-RR	GenBank Accession Number: NM_007425.3	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 11596	CloneNo.: 240774F11
Source: Rabbit	UNIPROT ID: Q62151-1	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000
Isotype: IgG	Full Name: advanced glycosylation end product-specific receptor	
	Calculated MW: 43 kDa	
	Observed MW: 42-55 kDa	

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: rat lung tissue, mouse lung tissue

IHC: mouse lung tissue,

Background Information

The receptor for advanced glycation end products (RAGE) is a cell surface transmembrane multiligand receptor, also known as Ager, is a multiligand member of the immunoglobulin superfamily. It has many isoforms and is predominantly expressed in the lung. AGER expression is upregulated in many diseases such as Alzheimer's disease, diabetes mellitus, atherosclerosis, and rheumatoid arthritis (PMID: 35099614).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83742-5-RR (RAGE antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 83742-5-RR (Ager/Rage antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Bi-layer interferometry (BLI) kinetic assays of 83608-4-RR against Human ATM were performed. The affinity constant is 2.58 nM. Bi-layer interferometry (BLI) kinetic assays of 83608-4-RR against Mouse RAGE were performed. The affinity constant is 6.41 nM.