

For Research Use Only

TNC/Tenascin-C Recombinant antibody



Catalog Number: 83767-1-RR

Basic Information

Catalog Number: 83767-1-RR	GenBank Accession Number: BC151843	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 3371	CloneNo.: 240844A2
Source: Rabbit	UNIPROT ID: P24821	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: tenascin C	
Immunogen Catalog Number: AG27122	Calculated MW: 2201 aa, 241 kDa	
	Observed MW: 220-350 kDa, 190-240 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : U-87 MG cells,
Species Specificity: human	

Background Information

Tenascin-C (TNC) is a large hexameric extracellular matrix glycoprotein of the tenascin family (PMID: 21818551). It is a multimodular protein containing multiple epidermal growth factor (EGF)-like repeats and fibronectin type III (FN III) domains (PMID: 1719530). Tenascin-C is highly expressed during embryonic development, particularly in the developing central nervous system, around motile cells and at epithelial-mesenchymal interaction sites (PMID: 25738825). In adult tissues, the expression and the distribution of TNC are typically limited under normal physiological conditions. It is upregulated during injury, inflammation, regeneration, and cancer (PMID: 7694605; 25120494). Tenascin-C is a diverse protein that can produce different functions including regulating cell adhesion, migration and proliferation.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



U-87 MG cells were subjected to SDS PAGE followed by western blot with 83767-1-RR (TNC antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 83767-1-RR against Human TNC/Tenascin-C were performed. The affinity constant is below 1 pM.