For Research Use Only

HDAC10 Recombinant antibody

Catalog Number:83788-4-RR



Basic Information	Catalog Number: 83788-4-RR	GenBank Accession Number: BC 125083	Purification Method: Protein A purfication	
	Size: 1000 µg/ml	Genel D (NCBI): 83933	CloneNo.: 240877E1	
	Source: Rabbit	UNIPROT ID: Q96958	Recommended Dilutions: WB 1:5000-1:50000	
	lsotype: IgG	Full Name: histone deacetylase 10		
	Immunogen Catalog Number: AG18851	Calculated MW: 669 aa, 71 kDa Observed MW: 70-75 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, ELISA Species Specificity: human	WB : Jurka	at cells,	
Background Informat	levels in liver and kidney. HDAC1 (DAC) and a catalytically inactive homologous recombination, prom	HDAC 10, also named as Histone deacetylase 10, is a 669 amino acid protein, which is widely expressed with high levels in liver and kidney. HDAC 10, which belongs to the class IIb subfamily, has an active deacetylase domain (DAC) and a catalytically inactive leucine-rich domain (LRD). HDAC 10 is determined that it plays a role in homologous recombination, promotes autophagy and survival in neuroblastoma cells, suppresses cervical cancer metastasis and facilitates the cell cycle.		
Storage	Storage Buffer:	Store at -20°C. Stable for one year after shipment. Storage Buffer:		
	PBS with 0.02% sodium azide and	PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20 [°] C storage		

 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data

 $\begin{array}{c} 180 \text{ kDa} \rightarrow \\ 140 \text{ kDa} \rightarrow \\ 100 \text{ kDa} \rightarrow \\ 75 \text{ kDa} \rightarrow \end{array}$

 $\begin{array}{l} 60 \text{ kDa} \rightarrow \\ 45 \text{ kDa} \rightarrow \end{array}$

 $35 \text{ kDa} \rightarrow$ $25 \text{ kDa} \rightarrow$ $15 \text{ kDa} \rightarrow$

10 kDa-



Jurkat cells were subjected to SDS PAGE followed by western blot with 83788-4-RR (HDAC10 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. Biolayer interferometry (BLI) kinetic assays of 83788-4-RR against Human HDAC 10 were performed. The affinity constant is 0.476 nM.