## For Research Use Only

## CXCL8/IL-8 Recombinant antibody

Catalog Number:83903-4-RR



**Basic Information** 

Catalog Number:

83903-4-RR

Size: 1000 ug/ml

Source: Rabbit

Isotype: IgG GenBank Accession Number:

BC013615 GeneID (NCBI):

3576

ENSEMBL Gene ID: ENSG00000169429

UNIPROT ID: P10145

Full Name: interleukin 8 Calculated MW: 99 aa, 11 kDa Observed MW: 11 kDa Purification Method:

Protein A purfication

CloneNo.: 240889C4

Recommended Dilutions:

WB 1:5000-1:50000

**Applications** 

**Tested Applications:** 

WB, ELISA

Species Specificity:

human

Positive Controls:

WB: PMA, LPS and Brefeldin A treated THP-1 cells,

## **Background Information**

Interleukin 8 (IL-8), also known as CXCL8, is a CXC chemokine family member. This chemokine is secreted by a variety of cell types including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, and various tumor cell lines in response to inflammatory stimuli. IL-8 has two primary functions. It induces chemotaxis in target cells, primarily neutrophils but also other granulocytes, causing them to migrate toward the site of infection. IL-8 also induces phagocytosis once they have arrived. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. IL-8 is also known to be a potent promoter of angiogenesis. IL-8 has been associated with tumor angiogenesis, metastasis, and poor prognosis in breast cancer. IL-8 may present a novel therapeutic target for estrogen driven breast carcinogenesis and tumor progression. The human IL-8 cDNA sequence predicts a protein of 99 amino acids. Removal of a 22-residue signal peptide generates a mature protein of 77 amino acids (~ 8 kDa).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.


Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data





K<sub>D</sub>=1.65×10<sup>-10</sup> K<sub>off</sub>=1.13×10<sup>-3</sup> K<sub>on</sub>=6.86×10<sup>6</sup>

PMA, LPS and Brefeldin A treated THP-1 cells were subjected to SDS PAGE followed by western blot with 83903-4-RR (CXCL8,IL8 antibody) at dilution of 1:10000 incubated at room temperature for 1.5

Biolayer interferometry (BLL) kinetic assays of 83903-4-RR against Human CXCL8/IL-8 were performed. The affinity constant is 0.165 nM.