For Research Use Only

## FANCD2 Recombinant antibody

Catalog Number:83946-4-RR





Basic Information	Catalog Number: 83946-4-RR	GenBank Accession Number: NM_001018115	Purification Method: Protein A purfication
	Size: 1000 µg/ml	GeneID (NCBI): 2177	CloneNo.: 241048G8
	Source: Rabbit	UNIPROT ID: Q9BXW9	Recommended Dilutions: WB 1:5000-1:50000
	Isotype: IgG Immunogen Catalog Number: AG29460	Full Name: Fanconi anemia, complementation group D2	
		Calculated MW: 164 kDa	
		Observed MW: 150-160 kDa	
Applications	Tested Applications: WB, FC (Intra), ELISA Species Specificity: human	Positive	e Controls:
		WB : HeLa cells, K-562 cells, MCF-7 cells, Jurkat cells U2OS cells	
Background Information	Amongst the FA genes, Fanconi Anemia Group D2 Protein (FANCD2) is the most evolutionarily conserved gene, demonstrating a high sequence similarity from lower eukaryotes to humans. Upon activation, Fanconi anemia group D2 protein (FANCD2) is monoubiquitinated at K561 and works in cooperation with other FA and non-FA proteins to repair DNA damage. In regards to the FA pathway, monoubiquitinated or activated FANCD2 is often portrayed as the functional representative for the functions of activated FA signaling. (PMID: 28825622)		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





LS K<sub>0</sub><100-10<sup>-12</sup> K<sub>00</sub><100-10<sup>-12</sup> K<sub>00</sub><100<sup>-10</sup> K<sub>00</sub><100<sup>-10</sup> 1.0 1.0 1.0 1.0 1.0 1.0 1.25 mM 2.25 mM 1.25 mM 0.25 mM 0.05 mM 0.25 mM 

Various lysates were subjected to SDS PAGE followed by western blot with 83946-4-RR (FANCD2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. 1x10^6 HeLa cells were intracellularly stained with 0.25 ug FANCD2 Recombinant antibody (83946-4-RR, Clone:241048G8) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+1)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.

