For Research Use Only

CD40 Recombinant antibody

Catalog Number:83972-5-RR



Basic Information

Catalog Number: 83972-5-RR

Size: 1000 µ g/ml Source: Rabbit Isotype: GenBank Accession Number: NM_011611.2 GeneID (NCBI): 21939 UNIPROT ID: P27512-1 Full Name: CD40 antigen

Calculated MW: 32 kDa Observed MW: 40-45 kDa Purification Method: Protein A purfication

CloneNo.: 240874B2

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, ELISA Species Specificity: mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: RAW 264.7 cells, J774A.1 cells IHC: mouse spleen tissue,

Background Information

CD40, also named as TNFRSF5 and Bp50, is a receptor for TNFSF5/CD40LG. It is a member of the TNFR superfamily and is responsible for the efficient activation of cell-mediated and humoral immune responses via activated T cells and B cells respectively. Defects in CD40 lead to hyper-IgM immunodeficiency syndrome type 3 (HIGM3) (PMID: 11675497). This is an autosomal recessive disorder that includes the inability of B cells to undergo isotype switching, a key step in the final differentiation of the humoral immune response, and an inability to mount an antibody-specific immune response.

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83972-5-RR (CD40 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 83972-5-RR (CD40 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse spleen tissue slide using 83972-5-RR (CD40 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 83972-5-RR against Mouse CD40 were performed. The affinity constant is 4.50 nM.