For Research Use Only

OGG1 Recombinant antibody


Catalog Number:83973-4-RR




Catalog Number: 83973-4-RR	GenBank Accession Number: BC000657	Purification Method: Protein A purfication
Size: 1000 µg/ml	GenelD (NCBI): 4968	CloneNo.: 240940D4
Source: Rabbit	UNIPROT ID: O 15527	Recommended Dilutions: IF/ICC 1:200-1:800
lsotype: IgG	Full Name: 8-oxoguanine DNA glycosylase	
Immunogen Catalog Number: AG7204	Calculated MW: 22 kDa, 36-40 kDa, 45-57 kDa	
Tested Applications: IF/ICC, FC (Intra), ELISA	Positive Controls:	
Species Specificity: human	IF/ICC : H	epuz cells,
The DNA damages induced by ROS contain base modification, base loss, and DNA single strand breaks, which are usually repaired by the base excision repair (BER) pathway in both prokaryotes and eukaryotes. OGG1 (The humar 8-oxoguanine glycosylase 1) is the primary enzyme in BER pathway, responsible for the excision of 7, 8-dihydro-8 oxoguanine (8-oxoG), a mutagenic base byproduct that occurs as a result of exposure to reactive oxygen species. There's 8 isoforms of OGG1, with calculated MW 22 kDa, 36-40 kDa and 45-57 kDa. The difference among these isoforms is the C-terminal (317-345aa).		
Storage: Store at -20°C. Stable for one year Storage Buffer:	after shipment.	
	 83973-4-RR Size: 1000 μ g/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG7204 Tested Applications: IF/ICC, FC (Intra), ELISA Species Specificity: human The DNA damages induced by RO usually repaired by the base excis	83973-4-RR BC000657 Size: GeneID (NCBI): 1000 µ g/ml 4968 Source: UNIPROT ID: Rabbit 015527 Isotype: Full Name: IgG 8-oxoguanine DNA glycosylase Immunogen Catalog Number: Calculated MW: AG7204 22 kDa, 36-40 kDa, 45-57 kDa Tested Applications: Positive C IF/ICC, FC (Intra), ELISA IF/ICC : H Species Specificity: IF/ICC : H human The DNA damages induced by ROS contain base modification, base loss: usually repaired by the base excision repair (BER) pathway in both profiles 8-oxoguanine (8-oxoG), a mutagenic base byproduct that occurs as a resu There's 8 isoforms of OGG1, with calculated MW 22 kDa, 36-40 kDa and isoforms is the C-terminal (317-345aa).

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using OGG1 antibody (83973-4-RR, Clone: 240940D4) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug Ogg1 Recombinant Antibody (83973-4-RR, Clone:240940D4) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.