## For Research Use Only

## NFKB2,p100 Recombinant antibody

Catalog Number:84022-6-RR



**Purification Method:** 

Protein A purfication

Recommended Dilutions:

WB 1:5000-1:50000 IF/ICC 1:125-1:500

CloneNo.:

241126F1

**Basic Information** 

Catalog Number: 84022-6-RR

Size: 1000 µg/ml

Source:

BC002844 GeneID (NCBI):

GenBank Accession Number:

4791 ENSEMBL Gene ID:

Rabbit ENSG00000077150 **UNIPROT ID:** Isotype: Q00653 Full Name:

Immunogen Catalog Number:

AG0662

nuclear factor of kappa light polypeptide gene enhancer in B-cells

2 (p49/p100) Calculated MW: 97 kDa Observed MW: 100-110 kDa

**Applications** 

**Tested Applications:** WB, IF/ICC, ELISA Species Specificity:

human

Positive Controls:

WB: HeLa cells, K-562 cells, Jurkat cells, SKOV-3 cells,

HL-60 cells

IF/ICC: U2OS cells,

## **Background Information**

NF-kappa-B is a pleiotropic transcription factor which is present in almost all cell types and is involved in many  $biological\ processed\ such\ as\ inflammation, immunity, differentiation, cell\ growth, tumorigenesis\ and\ apoptosis.\ NF-tumorigenesis\ and\ apoptosis\ processed\ such\ as\ inflammation, immunity, differentiation, cell\ growth, tumorigenesis\ and\ apoptosis\ processed\ such\ as\ inflammation, immunity, differentiation, cell\ growth, tumorigenesis\ and\ apoptosis\ processed\ such\ as\ inflammation, immunity, differentiation, cell\ growth, tumorigenesis\ and\ apoptosis\ processed\ such\ as\ inflammation\ processed\ such\ as\ inflamma$ kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. NFKB2 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p100 and generation of p52 by a cotranslational processing. The proteasomemediated process ensures the production of both p52 and p100 and preserves their independent function. P52 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. P52 and p100 are respectively the minor and major form; the processing of p100 being relatively poor. Isoform p49 is a subunit of the NF-kappa-B protein complex, which stimulates the HIV enhancer in synergy with p65. This antibody can bind p100 isoform of NFKB2 spefically, but not p52 isoform.

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 84022-6-RR (NFKB2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using NFKB2 antibody (84022-6-RR, Clone: 241126F1) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Biolayer interferometry (BLI) kinetic assays of 84022-6-RR against Human NFKB2,p100 were performed. The affinity constant is below 1 pM.