For Research Use Only

DSG1 Recombinant antibody

Catalog Number:84053-6-RR



Basic Information

Catalog Number: GenBank Accession Number: 84053-6-RR BC153001

 Size:
 GeneID (NCBI):

 1000 μ g/ml
 1828

Source: UNIPROT ID:
Rabbit Q02413
Isotype: Full Name:
IgG desmoglein 1
Immunogen Catalog Number: Calculated MW:
AG20184 1049 aa, 114 kDa

Observed MW: 160 kDa Purification Method: Protein A purfication

CloneNo.: 241093G11

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000

Applications

Tested Applications: WB, IHC, ELISA Species Specificity:

human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 **Positive Controls:**

WB: A431 cells, HEK-293 cells, HeLa cells, rat skin

tissue

IHC: mouse skin tissue,

Background Information

Desmosomes are cell-cell junctions between epithelial, myocardial, and certain other cell types. Desmosomal cadherins, consisting of four desmogleins (DSG1-4) and three desmocollins (DSC1-3) in humans, mediate adhesion through calcium-dependent homophilic/heterophilic interactions. DSG1 is a single-pass transmembrane glycoprotein highly expressed in the epidermis and localized primarily within the suprabasal epithelial layers (PMID: 16286477; 24220297). DSG1 mediates intercellular adhesion and is crucial in maintaining epidermal integrity and barrier function (PMID: 23974871). It is also involved in epithelial cell differentiation (PMID: 23524961). Mutations in the DSG1 gene can cause the autosomal dominant disorder to striate palmoplantar keratoderma and a syndrome featuring severe dermatitis, multiple allergies, and metabolic wasting (SAM syndrome) (PMID: 29315490; 23974871).

Storage


Storage

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84053-6-RR (DSG1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse skin tissue slide using 84053-6-RR (DSG1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse skin tissue slide using 84053-6-RR (DSG1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 84053-6-RR against Human DSG1 were performed. The affinity constant is 71.2 pM.